

| | Albania | Cyprus | Egypt | Greece | Italy | Tunisia |
|--|---------------------|--------------------|-------------------|--------------------|--------------------|-------------------|
| Study sites | 1 | 1 | 1 | 1 | 11 | 1 |
| Start of recruitment | Sept 2013 | Mar 2015 | Apr 2013 | Nov 2012 | May 2013 | Feb 2015 |
| Enrolled subjects | 5 | 13 | 154 | 23 | 101 | 14 |
| With complete records | 5 | 13 | 148 | 23 | 97 | 11 |
| Age (median, range, in years) ¹ | 13.3 (10.3-14.4) | 11.1 (5.0-14.1) | 4.6 (0.6-17.6) | 10.5 (3.3-16.9) | 10.5 (1.1-17.4) | 7.5 (4.0-12.6) |
| Follow-up (total person-years) | 13.7 | 30.8 | 253.6 | 73.6 | 326.2 | 19.3 |
| Retrospective observation | 11.3 | 29.4 | 230.9 | 70.3 | 302.6 | 19.3 |
| Prospective observation | 2.4 | 1.5 | 22.7 | 3.4 | 23.6 | 0 |
| Per patient (median, IQR) | 3.3 (1.9-3.5) | 1.6 (0.7-3.7) | 1.2 (0.7-2.1) | 2.7 (1.4-4.8) | 2.7 (1.4-4.9) | 1.1 (0.3-3.4) |

¹ P-value <0.001 (Kruskal-Wallis test).

Table 4. DEEP-3 study characteristics