

**Table 2:** Patient and bereaved carer endpoints

<b>I. Measured by questionnaire</b>	<b>Measure</b>
<b>Primary endpoints:</b>	
- Quality of life	EORTC QLQ-C30 4-item emotional functioning scale EORTC emotional functioning short-form based on CAT item bank
- Symptoms	EORTC QLQ-C15-PAL
<b>Secondary endpoints:</b>	
- Shared decision-making	APECC
- Patient involvement	Self-constructed questions
- Satisfaction with care	EORTC IN-PATSAT32
- Coping with illness	COPE
- Satisfaction with the intervention	Self-constructed questions
- Socio demographic measures	Self-constructed questions
- Quality of end-of-life care	VOICES-SF*
- Bereaved carer wellbeing	HADS; IES*
<b>II. Obtained from medical files</b>	
- Survival; date and place of death (if applicable)	
- Completion and content of advance directives; preferences for care; assignment of proxy decision-maker; physician orders	
- Diagnostic procedures and treatments received by the patient, hospitalisations and specialist palliative care input.	
<b>III. Obtained from intervention sessions and qualitative interviews</b>	
Systematic cross-cultural comparison of patient experiences, responses and concerns.	

\* These endpoints are measured by the bereaved carer questionnaire and not by the patient questionnaire