Table 3. Sociodemographic and clinical characteristics of ACTION participants

Sociodemographic Characteristics	Control group (n = 675)	Intervention group (n = 442)	p-value
Age (years), mean (SD) Range Missing	66.3 (9.5) 30-91 <i>3</i>	66.3 (10.3) 18-89	0.92
Years of education, mean (SD)	12.9 (4.7)	13.1 (4.5)	0.42
Missing	94	56	
Female gender, n (%)	268 (40)	173 (39)	0.85
Living with a spouse, n (%)	497 (74)	303 (69)	0.16
Missing	17	10	2.2.5
Having children, n (%)	583 (86)	376 (85)	0.72
Missing	7	8	0.72
Religion, n (%)		0	0.37
Religious	341 (51)	207 (47)	0.57
Not religious	228 (34)	174 (39)	
Prefers not to specify	93 (14)	51 (12)	
Missing Considering oneself member of	13	10	
minority group, n (%)	7 (1)	3 (1)	0.92
Missing	26	18	
Country of residence, n (%)	<u> </u>		<0.001
Belgium	135 (20)	72 (16)	
Denmark	68 (10)	68 (15)	
Italy	139 (21)	31 (7)	
The Netherlands	168 (25)	84 (19)	
Slovenia	25 (4)	72 (16)	
The United Kingdom	140 (21)	115 (26)	
Clinical Characteristic	110 (21)	115 (25)	
Diagnosis, n (%)			<0.001
Lung cancer, stage III or IV	341 (51)	273 (62)	
Colorectal cancer, stage IV	332 (49)	168 (38)	
Missing	2	1	
Years since diagnosis, mean (SD) Range	1.7 (2.4) 0.1 – 33.3	1.2 (1.7) 0.1 – 11.5	<0.001
Missing	1	7	
Years since diagnosis of current			
stage, mean (SD) Range	1.0 (1.4) 0.1 – 11.1	0.6 (0.9) 0.1 – 5.8	<0.001
Missing	6	10	
Receiving systemic treatment ¹ , n (%)	595 (89)	349 (79)	<0.001
Missing	3	2	
WHO performance status ² , n (%)			<0.001

3	8 (1)	10 (2)	
2	55 (8)	74 (17)	
1	343 (51)	243 (55)	
0	261 (39)	109 (25)	
Missing	8	6	

¹Includes chemotherapy, immunotherapy, and targeted therapy.